

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 December 2001 (27.12.2001)

PCT

(10) International Publication Number
WO 01/97788 A2

- (51) International Patent Classification⁷: **A61K 31/00**
- (21) International Application Number: **PCT/EP01/06850**
- (22) International Filing Date: **18 June 2001 (18.06.2001)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
09/597,135 20 June 2000 (20.06.2000) US
60/267,689 9 February 2001 (09.02.2001) US
- (71) Applicant (for all designated States except AT, US): **NOVARTIS AG [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).**
- (71) Applicant (for AT only): **NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT M.B.H. [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).**
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **HOROWITZ, Zebulun, D. [US/US]; 83 Blackburn Road, Basking Ridge, NJ 07920 (US). RICHARDSON, Peter, C. [GB/US]; 4 Spring Lake Drive, Far Hills, NJ 07931 (US). TRECHSEL, Ulrich [CH/CH]; Grosshaus, CH-7242 Luzern (CH).**
- (74) Agent: **BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Department, CH-4002 Basel (CH).**
- (81) Designated States (national): **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.**
- (84) Designated States (regional): **ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).**
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/97788 A2

(54) Title: **METHOD OF ADMINISTERING BISPHOSPHONATES**

(57) Abstract: Bisphosphonates, in particular more potent N-bisphosphonates such as zoledronic acid and derivatives, can be used with satisfactory results for prolonged inhibition of bone resorption in conditions of abnormally increased bone turnover, e.g. osteoporosis, by intermittent administration, wherein the periods between bisphosphonate administrations are longer than was previously considered appropriate, e.g. a dosing interval of at least about 6 months or less frequently.

METHOD OF ADMINISTERING BIPHOSPHONATES

This invention relates to bisphosphonates, in particular to the pharmaceutical use of bisphosphonates in the treatment of conditions of abnormally increased bone turnover, such as osteoporosis:

Bisphosphonates are widely used to inhibit osteoclast activity in a variety of both benign and malignant diseases in which bone resorption is increased. Thus bisphosphonates have recently become available for long-term treatment of patients with Multiple Myeloma (MM). These pyrophosphate analogs not only reduce the occurrence of skeletal related events but they also provide patients with clinical benefit and improve survival. Bisphosphonates are able to prevent bone resorption *in vivo*; the therapeutic efficacy of bisphosphonates has been demonstrated in the treatment of Paget's disease of bone, tumour-induced hypercalcemia and, more recently, bone metastasis and multiple myeloma (MM) (for review see Fleisch H 1997 Bisphosphonates clinical. In Bisphosphonates in Bone Disease. From the Laboratory to the Patient. Eds: The Parthenon Publishing Group, New York/London pp 68-163). The mechanisms by which bisphosphonates inhibit bone resorption are still poorly understood and seem to vary according to the bisphosphonates studied. Bisphosphonates have been shown to bind strongly to the hydroxyapatite crystals of bone, to reduce bone turn-over and resorption, to decrease the levels of hydroxyproline or alkaline phosphatase in the blood, and in addition to inhibit both the activation and the activity of osteoclasts.

In addition bisphosphonates have been proposed for use in the treatment of osteoporosis. Thus for instance, as described in USP 4,812,304 (Procter & Gamble) a method has been proposed for treating or preventing osteoporosis in humans comprising administering to a subject afflicted with or at risk to osteoporosis a bone cell activating compound and a bone resorption inhibiting polyphosphonate according to a regime consisting of one or more cycles, whereby each cycle consists of: (a) a bone activating period of from about 1 day to about 5 days during which a bone cell activating amount of a bone cell activating compound is administered to said subject; followed by (b) a bone resorption inhibition period of from about 10 days to about 20 days during which ethane-1-

hydroxy-1,1-diphosphonic acid, or a pharmaceutically acceptable salt or ester thereof, is administered daily to said subject in an amount from about 0.25 mgP/kg/day to about 3.3mgP/kg/day; followed by (c) a rest period of from about 70 days to about 180 days during which the subject receives neither a bone cell activating compound nor a bone resorption inhibiting polyphosphonate.

Also, for example, USP 4,761,406 (Procter & Gamble) proposes a method for treating osteoporosis, in humans or lower animals afflicted with or at risk of osteoporosis, comprising administering to said human or lower animal an effective amount of a bone resorption inhibiting polyphosphonate according to the following schedule: (a) a period of from about 1 day to about 90 days during which said bone resorption inhibiting polyphosphonate is administered daily in a limited amount, followed by (b) a rest period of from about 50 days to about 120 days, and (c) repeating (a) and (b) two or more times where a net increase in bone mass of said human or animal results.

Surprisingly we have now found that bisphosphonates, in particular more potent nitrogen-containing bisphosphonates, can be used for prolonged inhibition of bone resorption in conditions of abnormally increased bone turnover by intermittent administration, wherein the periods between bisphosphonate administrations are longer than was previously considered appropriate to achieve satisfactory treatment. In particular and contrary to expectation we have found that satisfactory treatment results can be obtained even when the dosing intervals greatly exceed the natural bone remodelling cycle.

Accordingly the present invention provides a method for the treatment of conditions of abnormally increased bone turnover in a patient in need of such treatment which comprises intermittently administering an effective amount of a bisphosphonate to the patient, wherein the period between administrations of bisphosphonate is at least about 6 months.

The invention further provides use of a bisphosphonate in the preparation of a medicament for the treatment of conditions of abnormally increased bone turnover in which the bisphosphonate is

administered intermittently and in which the period between administrations of bisphosphonate is at least about 6 months.


Conditions of abnormally increased bone turnover which may be treated in accordance with the present invention include: treatment of postmenopausal osteoporosis, e.g. to reduce the risk of osteoporotic fractures; prevention of postmenopausal osteoporosis, e.g. prevention of postmenopausal bone loss; treatment or prevention of male osteoporosis; treatment or prevention of corticosteroid-induced osteoporosis and other forms of bone loss secondary to or due to medication, e.g. diphenylhydantoin, thyroid hormone replacement therapy; treatment or prevention of bone loss associated with immobilisation and space flight; treatment or prevention of bone loss associated with rheumatoid arthritis, osteogenesis imperfecta, hyperthyroidism, anorexia nervosa, organ transplantation, joint prosthesis loosening, and other medical conditions. For example, such other medical conditions may include treatment or prevention of periarticular bone erosions in rheumatoid arthritis; treatment of osteoarthritis, e.g. prevention/treatment of subchondral osteosclerosis, subchondral bone cysts, osteophyte formation, and of osteoarthritic pain, e.g. by reduction in intra-osseous pressure; treatment or prevention of hypercalcemia resulting from excessive bone resorption secondary to hyperparathyroidism, thyrotoxicosis, sarcoidosis or hypervitaminosis D.

Thus in the present description the terms "treatment" or "treat" refer to both prophylactic or preventative treatment as well as curative or disease modifying treatment, including treatment of patients at risk of contracting the disease or suspected to have contracted the disease as well as patients who are ill or have been diagnosed as suffering from a disease or medical condition. In particularly preferred embodiments the invention may be used for the prophylactic treatment of osteoporosis and similar diseases. Thus for example, bisphosphonate may be administered to individuals at risk of developing osteoporosis on a regular basis at dosing intervals of at least about 6 months, e.g. bisphosphonates may be administered routinely to postmenopausal women at dosing intervals of once every 6 months or less frequently, e.g. annually.

In accordance with the present invention the bisphosphonate dosing interval is at least about 6 months, e.g. once every 180 days, or less frequently, conveniently once a year, or at any interval in

between, e.g. once every 7, 8, 9, 10, or 11 months. Dosing intervals of greater than once per year may be used, e.g. about once every 18 months or about once every 2 years, or even less frequently, e.g. a frequency of up to about once every 3 years or less often.

The bisphosphonates used in the present invention are typically those which inhibit bone resorption. Such compounds characteristically contain two phosphonate groups attached to a single carbon atom, forming a "P-C-P" structure, e.g. in a compound of formula I



wherein

X is hydrogen, hydroxyl, amino, alkanoyl, or an amino group mono- or disubstituted by C₁-C₄ alkyl;

R is hydrogen or C₁-C₄ alkyl and


Rx is an optionally substituted hydrocarbyl group;

and pharmaceutically acceptable salts thereof or any hydrate thereof.

Thus, for example, suitable bisphosphonates for use in the invention may include the following compounds or a pharmaceutically acceptable salt thereof, or any hydrate thereof: 3-amino-1-hydroxypropane-1,1-diphosphonic acid (pamidronic acid), e.g. pamidronate (APD); 3-(N,N-dimethyl-amino)-1-hydroxypropane-1,1-diphosphonic acid, e.g. dimethyl-APD; 4-amino-1-hydroxybutane-1,1-diphosphonic acid (alendronic acid), e.g. alendronate; 1-hydroxy-ethidene-bisphosphonic acid, e.g. etidronate; 1-hydroxy-3-(methylpentylamino)-propylidene-bisphosphonic acid, (ibandronic acid), e.g. ibandronate; 6-amino-1-hydroxyhexane-1,1-diphosphonic acid, e.g. amino-hexyl-BP; 3-(N-methyl-N-pentylamino)-1-hydroxypropane-1,1-diphosphonic acid, e.g. methyl-pentyl-APD (= BM 21.0955); 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid, e.g. zoledronic acid; 1-hydroxy-2-(3-pyridyl)ethane-1,1-diphosphonic acid (risedronic acid), e.g. risedronate, including N-methyl

pyridinium salts thereof, for example N-methyl pyridinium iodides such as NE-10244 or NE-10446; 1-(4-chlorophenylthio)methane-1,1-diphosphonic acid (tiludronic acid), e.g. tiludronate; 3-[N-(2-phenylthioethyl)-N-methylamino]-1-hydroxypropane-1,1-diphosphonic acid; 1-hydroxy-3-(pyrrolidin-1-yl)propane-1,1-diphosphonic acid, e.g. EB 1053 (Leo); 1-(N-phenylaminothiocarbonyl)methane-1,1-diphosphonic acid, e.g. FR 78844 (Fujisawa); 5-benzoyl-3,4-dihydro-2H-pyrazole-3,3-diphosphonic acid tetraethyl ester, e.g. U-81581 (Upjohn); 1-hydroxy-2-(imidazo[1,2-a]pyridin-3-yl)ethane-1,1-diphosphonic acid, e.g. YM 529; and 1,1-dichloromethane-1,1-diphosphonic acid (clodronic acid), e.g. clodronate; YM175.

Preferred bisphosphonates for use in the present invention are N-bisphosphonates, i.e. compounds which in addition to the characteristic geminal bisphosphonates moiety (e.g. "P-C-P") comprise a nitrogen-containing side chain, e.g. a compound of formula I'



wherein

X is hydrogen, hydroxyl, amino, alkanoyl, or an amino group mono- or disubstituted by C₁-C₄ alkyl;

R is hydrogen or C₁-C₄ alkyl and


Rx' is a side chain which contains an optionally substituted amino group, or a nitrogen containing heterocycle (including aromatic nitrogen-containing heterocycles),

and pharmaceutically acceptable salts thereof or any hydrate thereof.

Thus, for example, suitable N-bisphosphonates for use in the invention may include the following compounds or a pharmaceutically acceptable salt thereof, or any hydrate thereof: 3-amino-1-hydroxypropane-1,1-diphosphonic acid (pamidronic acid), e.g. pamidronate (APD); 3-(N,N-dimethylamino)-1-hydroxypropane-1,1-diphosphonic acid, e.g. dimethyl-APD; 4-amino-1-

hydroxybutane-1,1-diphosphonic acid (alendronic acid), e.g. alendronate; 1-hydroxy-3-(methylpentylamino)-propylidene-bisphosphonic acid, ibandronic acid, e.g. ibandronate; 6-amino-1-hydroxyhexane-1,1-diphosphonic acid, e.g. amino-hexyl-BP; 3-(N-methyl-N-n-pentylamino)-1-hydroxypropane-1,1-diphosphonic acid, e.g. methyl-pentyl-APD (= BM 21.0955); 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid, e.g. zoledronic acid; 1-hydroxy-2-(3-pyridyl)ethane-1,1-diphosphonic acid (risedronic acid), e.g. risedronate, including N-methyl pyridinium salts thereof, for example N-methyl pyridinium iodides such as NE-10244 or NE-10446; 3-[N-(2-phenylthioethyl)-N-methylamino]-1-hydroxypropane-1,1-diphosphonic acid; 1-hydroxy-3-(pyrrolidin-1-yl)propane-1,1-diphosphonic acid, e.g. EB 1053 (Leo); 1-(N-phenylaminothiocarbonyl)methane-1,1-diphosphonic acid, e.g. FR 78844 (Fujisawa); 5-benzoyl-3,4-dihydro-2H-pyrazole-3,3-diphosphonic acid tetraethyl ester, e.g. U-81581 (Upjohn); and 1-hydroxy-2-(imidazo[1,2-a]pyridin-3-yl)ethane-1,1-diphosphonic acid, e.g. YM 529.

In one embodiment a particularly preferred N-bisphosphonate for use in the invention comprises a compound of Formula II



wherein

Het is an imidazole, oxazole, isoxazole, oxadiazole, thiazole, thiadiazole, pyridine, 1,2,3-triazole, 1,2,4-triazole or benzimidazole radical, which is optionally substituted by alkyl, alkoxy, halogen, hydroxyl, carboxyl, an amino group optionally substituted by alkyl or alkanoyl radicals or a benzyl radical optionally substituted by alkyl, nitro, amino or aminoalkyl;

A is a straight-chained or branched, saturated or unsaturated hydrocarbon moiety containing from 1 to 8 carbon atoms;


X' is a hydrogen atom, optionally substituted by alkanoyl, or an amino group optionally

substituted by alkyl or alkanoyl radicals, and

R is a hydrogen atom or an alkyl radical,

and the pharmacologically acceptable salts thereof.

In a further embodiment a particularly preferred bisphosphonate for use in the invention comprises a compound of Formula III



wherein

Het' is a substituted or unsubstituted heteroaromatic five-membered ring selected from the group consisting of imidazolyl, imidazolyl, isoxazolyl, oxazolyl, oxazolyl, thiazolyl, thiazolyl, triazolyl, oxadiazolyl and thiadiazolyl wherein said ring can be partly hydrogenated and wherein said substituents are selected from at least one of the group consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, phenyl, cyclohexyl, cyclohexylmethyl, halogen and amino and wherein two adjacent alkyl substituents of Het can together form a second ring;


Y is hydrogen or C₁-C₄ alkyl;

X'' is hydrogen, hydroxyl, amino, or an amino group substituted by C₁-C₄ alkyl, and

R is hydrogen or C₁-C₄ alkyl;

as well as the pharmacologically acceptable salts and isomers thereof.

In a yet further embodiment a particularly preferred bisphosphonate for use in the invention comprises a compound of Formula IV



wherein

Het''' is an imidazolyl, 2H-1,2,3-, 1H-1,2,4- or 4H-1,2,4-triazolyl, tetrazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl or thiadiazolyl radical which is unsubstituted or C-mono- or di-substituted by lower alkyl, by lower alkoxy, by phenyl which may in turn be mono- or disubstituted by lower alkyl, lower alkoxy and/or halogen, by hydroxy, by di-lower alkylamino, by lower alkylthio and/or by halogen and is N-substituted at a substitutable N-atom by lower alkyl or by phenyl-lower alkyl which may in turn be mono- or di-substituted in the phenyl moiety by lower alkyl, lower alkoxy and/or halogen, and

R₂ is hydrogen, hydroxy, amino, lower alkylthio or halogen,

lower radicals having up to and including 7 C-atoms,

or a pharmacologically acceptable salt thereof.

Examples of particularly preferred N-bisphosphonates for use in the invention are:

2-(1-Methylimidazol-2-yl)-1-hydroxyethane-1,1-diphosphonic acid;

2-(1-Benzylimidazol-2-yl)-1-hydroxyethane-1,1-diphosphonic acid;

2-(1-Methylimidazol-4-yl)-1-hydroxyethane-1,1-diphosphonic acid;

1-Amino-2-(1-methylimidazol-4-yl)ethane-1,1-diphosphonic acid;

1-Amino-2-(1-benzylimidazol-4-yl)ethane-1,1-diphosphonic acid;

2-(1-Methylimidazol-2-yl)ethane-1,1-diphosphonic acid;

2-(1-Benzylimidazol-2-yl)ethane-1,1-diphosphonic acid;

2-(Imidazol-1-yl)-1-hydroxyethane-1,1-diphosphonic acid;

2-(Imidazol-1-yl)ethane-1,1-diphosphonic acid;

2-(4H-1,2,4-triazol-4-yl)-1-hydroxyethane-1,1-diphosphonic acid;

2-(Thiazol-2-yl)ethane-1,1-diphosphonic acid;
2-(Imidazol-2-yl)ethane-1,1-diphosphonic acid;
2-(2-Methylimidazol-4(5)-yl)ethane-1,1-diphosphonic acid;
2-(2-Phenylimidazol-4(5)-yl)ethane-1,1-diphosphonic acid;
2-(4,5-Dimethylimidazol-1-yl)-1-hydroxyethane-1,1-diphosphonic acid, and
2-(2-Methylimidazol-4(5)-yl)-1-hydroxyethane-1,1-diphosphonic acid,
and pharmacologically acceptable salts thereof.

The most preferred N-bisphosphonate for use in the invention is 2-(imidazol-1-yl)-1-hydroxyethane-1,1-diphosphonic acid (zoledronic acid) or a pharmacologically acceptable salt thereof.

Pharmacologically acceptable salts are preferably salts with bases, conveniently metal salts derived from groups Ia, Ib, IIa and IIb of the Periodic Table of the Elements, including alkali metal salts, e.g. potassium and especially sodium salts, or alkaline earth metal salts, preferably calcium or magnesium salts, and also ammonium salts with ammonia or organic amines.

Especially preferred pharmaceutically acceptable salts are those where one, two, three or four, in particular one or two, of the acidic hydrogens of the bisphosphonic acid are replaced by a pharmaceutically acceptable cation, in particular sodium, potassium or ammonium, in first instance sodium.

A very preferred group of pharmaceutically acceptable salts is characterized by having one acidic hydrogen and one pharmaceutically acceptable cation, especially sodium, in each of the phosphonic acid groups.

The bisphosphonic acid derivatives specifically mentioned above are well known from the literature. This includes their manufacture (see e.g. EP-A-513760, pp. 13-48). For example, 3-amino-1-hydroxypropane-1,1-diphosphonic acid is prepared as described e.g. in US patent 3,962,432 as well

as the disodium salt as in US patents 4,639,338 and 4,711,880, and 1-hydroxy-2-(imidazol-1-yl)-ethane-1,1-diphosphonic acid is prepared as described e.g. in US patent 4,939,130.

The bisphosphonates (hereinafter referred to as the Agents of the Invention) may be used in the form of an isomer or of a mixture of isomers where appropriate, typically as optical isomers such as enantiomers or diastereoisomers or geometric isomers, typically cis-trans isomers. The optical isomers are obtained in the form of the pure antipodes and/or as racemates.

The Agents of the Invention can also be used in the form of their hydrates or include other solvents used for their crystallisation.

The Agents of the Invention are preferably used in the form of pharmaceutical compositions that contain a therapeutically effective amount of active ingredient optionally together with or in admixture with inorganic or organic, solid or liquid, pharmaceutically acceptable carriers which are suitable for administration,

The Agents of the Invention may be administered alone or in combination with other bone active drugs, either in fixed combinations or separately both physically and in time, including hormones, such as a steroid hormone, e.g. an estrogen; a partial estrogen agonist, or estrogen-gestagen combination; a calcitonin or an analogue or derivative thereof, e.g. salmon, eel or human calcitonin parathyroid hormone or analogues thereof, e.g. e.g. PTH (1-84), PTH (1-34), PTH (1-36), PTH (1-38), PTH (1-31)NH₂ or PTS 893; a SERM (Selective Estrogen Receptor Modulator) e.g. raloxifene, lasofoxifene, TSE-424, FC1271, Tibolone (Livial ®); vitamin D or an analog. Such additional bone active drugs may be administered more frequently than the bisphosphonate.

The pharmaceutical compositions may be, for example, compositions for enteral, such as oral, rectal, aerosol inhalation or nasal administration, compositions for parenteral, such as intravenous or subcutaneous administration, or compositions for transdermal administration (e.g. passive or iontophoretic).

Preferably, the pharmaceutical compositions are adapted to oral or parenteral (especially intravenous, subcutaneous, intramuscular, or transdermal) administration. Intravenous and oral, first and foremost, intravenous administration is considered to be of particular importance. Preferably the bisphosphonate active ingredient is in the form of a parenteral, most preferably an intravenous form.

The particular mode of administration and the dosage may be selected by the attending physician taking into account the particulars of the patient, especially age, weight, life style, activity level, hormonal status (e.g. post-menopausal) and bone mineral density as appropriate.

The dosage of the Agents of the Invention may depend on various factors, such as effectiveness and duration of action of the active ingredient, e.g. including the relative potency of the bisphosphonate used, mode of administration, warm-blooded species, and/or sex, age, weight and individual condition of the warm-blooded animal.

Normally the dosage is such that a single dose of the bisphosphonate active ingredient from 0.005 - 20 mg/kg, especially 0.01 - 10 mg/kg, is administered to a warm-blooded animal weighing approximately 75kg.

"mg/kg" means mg drug per kg body weight of the mammal - including man - to be treated.

The dose mentioned above is typically administered intermittently with a period of at least 6 months between doses. The period between bisphosphonate administrations may be longer, e.g. conveniently once per year, once per 18 months or once every 2 years, or even longer, or any period in between.

Formulations in single dose unit form contain preferably from about 1% to about 90%, and formulations not in single dose unit form contain preferably from about 0.1% to about 20%, of the active ingredient. Single dose unit forms such as ampoules of infusion solution or solid for preparation

of infusion solution doses, capsules, tablets or dragées contain e.g. from about 0.5 mg to about 500mg of the active ingredient. It will be appreciated that the actual unit dose used will depend upon the potency of the bisphosphonates, the dosing interval and route of administration amongst other things. Thus the size of the unit dose is typically lower for more potent bisphosphonates and greater the longer the dosing interval. For example, for more potent, N-bisphosphonates such as zoledronic acid a unit dose of from about 1 up to about 10 mg may be used for parenteral, e.g. intravenous, administration. For example, also for more potent N-bisphosphonates a unit dose of from about 1 to about 5 mg may be used parenterally for dosing once every 6 months; whereas a dose of from about 2 up to about 10 mg may be used for once a year parenteral dosing.

Unit doses may be administered as a single or divided dose, i.e. a dose in which the unit dose is divided into two or more equal or unequal parts and in which the parts are administered to the patient at the same time, during overlapping time periods or at separate time points. When the unit dose is administered as a divided dose at separate time points, the interval between the separate administrations of the divided dose may be from hours, e.g. 1 hour, up to about 1 month (approximately 30 days). In accordance with the invention, the time interval between administration of the last part of the divided dose and administration of the first part of the next, following divided dose is at least 6 months or longer, e.g. about 1 year.

Thus, for example, a 10 mg unit dose may be administered as two equal 5 mg parts at an interval of about 1 week to about 1 month, e.g. about 2 weeks, between administration of the parts. Alternatively, for example, a 5 mg unit dose may be administered as two unequal parts of 4 mg and 1 mg (or 3 mg and 2 mg) at an interval of from 1 to 3 days to 1 to 3 weeks, e.g. about 1 week, between administration of the parts.

Pharmaceutical preparations for enteral and parenteral administration are, for example, those in dosage unit forms, such as dragées, tablets or capsules and also ampoules. They are prepared in a manner known *per se*, for example by means of conventional mixing, granulating, confectioning, dissolving or lyophilising processes. For example, pharmaceutical preparations for oral administration

can be obtained by combining the active ingredient with solid carriers, where appropriate granulating a resulting mixture, and processing the mixture or granulate, if desired or necessary after the addition of suitable adjuncts, into tablets or dragée cores.

Suitable carriers are especially fillers, such as sugars, for example lactose, saccharose, mannitol or sorbitol, cellulose preparations, and also binders, such as starch pastes, using, for example, corn, wheat, rice or potato starch, gelatin, tragacanth, methylcellulose and/or polyvinylpyrrolidone and, if desired, disintegrators, such as the above-mentioned starches, also carboxymethyl starch, crosslinked polyvinylpyrrolidone, agar or alginic acid or a salt thereof, such as sodium alginate. Adjuncts are especially flow-regulating agents and lubricants, for example silicic acid, talc, stearic acid or salts thereof, such as magnesium or calcium stearate, and/or polyethylene glycol. Dragee cores are provided with suitable coatings that may be resistant to gastric juices, there being used, inter alia, concentrated sugar solutions that optionally contain gum arabic, talc, polyvinylpyrrolidone, polyethylene glycol and/or titanium dioxide, or lacquer solutions in suitable organic solvents or solvent mixtures or, to produce coatings that are resistant to gastric juices, solutions of suitable cellulose preparations, such as acetylcellulose phthalate or hydroxypropylmethylcellulose phthalate. Colouring substances or pigments may be added to the tablets or dragee coatings, for example for the purpose of identification or to indicate different doses of active ingredient.

Other orally administrable pharmaceutical preparations are dry-filled capsules made of gelatin, and also soft, sealed capsules made of gelatin and a plasticiser, such as glycerol or sorbitol. The dry-filled capsules may contain the active ingredient in the form of a granulate, for example in admixture with fillers, such as lactose, binders, such as starches, and/or glidants, such as talc or magnesium stearate, and, where appropriate, stabilisers. In soft capsules the active ingredient is preferably dissolved or suspended in suitable liquids, such as fatty oils, paraffin oil or liquid polyethylene glycols, it being possible also for stabilisers to be added.

Parenteral formulations are especially injectable fluids that are effective in various manners, such as, intramuscularly, intraperitoneally, intranasally, intradermally, subcutaneously or preferably

intravenously. Such fluids are preferably isotonic aqueous solutions or suspensions which can be prepared before use, for example from lyophilised preparations which contain the active ingredient alone or together with a pharmaceutically acceptable carrier, or from solution concentrates. The pharmaceutical preparations may be sterilised and/or contain adjuncts, for example preservatives, stabilisers, wetting agents and/or emulsifiers, solubilisers, salts for regulating the osmotic pressure and/or buffers.

Suitable formulations for transdermal application include an effective amount of the active ingredient with carrier. Advantageous carriers include absorbable pharmacologically acceptable solvents to assist passage through the skin of the host. Characteristically, transdermal devices are in the form of a bandage comprising a backing member, a reservoir containing the compound optionally with carriers, optionally a rate controlling barrier to deliver the active ingredient to the skin of the host at a controlled and predetermined rate over a prolonged period of time, and means to secure the device to the skin.

The following Examples illustrate the invention described hereinbefore.

In the following Examples the term "active ingredient" is to be understood as being any one of the bisphosphonic acid derivatives mentioned above as being useful according to the present invention.

EXAMPLES

Example 1: Capsules containing coated pellets of active ingredient; for example, disodium pamidronate pentahydrate, as active ingredient:

Core pellet:

active ingredient (ground)	197.3 mg
Microcrystalline cellulose (Avicel® PH 105)	52.7 mg
	<hr/>
	250.0 mg

+ Inner coating:

Cellulose HP-M 603	10.0 mg
Polyethylene glycol	2.0 mg
Talc	8.0 mg
	<hr/>
	270.0 mg

+ Gastric juice-resistant outer coating:

Eudragit® L 30 D (solid)	90.0 mg
Triethyl citrate	21.0 mg
Antifoam® AF	2.0 mg
Water	
Talc	7.0 mg
	<hr/>
	390.0 mg

A mixture of disodium pamidronate with Avicel® PH 105 is moistened with water and kneaded, extruded and formed into spheres. The dried pellets are then successively coated in the fluidized bed

with an inner coating, consisting of cellulose HP-M 603, polyethylene glycol (PEG) 8000 and talc, and the aqueous gastric juice-resistant coat, consisting of Eudragit® L 30 D, triethyl citrate and Antifoam® AF. The coated pellets are powdered with talc and filled into capsules (capsule size 0) by means of a commercial capsule filling machine, for example Höfliger and Karg.

Example 2: Monolith adhesive transdermal system, containing as active ingredient, for example, 1-hydroxy-2-(imidazol-1-yl)-ethane-1,1-diphosphonic acid:

Composition:

polyisobutylene (PIB) 300 (Oppanol B1, BASF)	5.0 g
PIB 35000 (Oppanol B10, BASF)	3.0 g
PIB 1200000 (Oppanol B100, BASF)	9.0 g
hydrogenated hydrocarbon resin (Escorez 5320, Exxon)	43.0 g
1-dodecylazacycloheptan-2-one (Azone, Nelson Res., Irvine/CA)	20.0 g
active ingredient	<u>20.0 g</u>
Total	100.0 g

Preparation:

The above components are together dissolved in 150 g of special boiling point petroleum fraction 100-125 by rolling on a roller gear bed. The solution is applied to a polyester film (Hostaphan, Kalle) by means of a spreading device using a 300mm doctor blade, giving a coating of about 75 g/m². After drying (15 minutes at 60°C), a silicone-treated polyester film (thickness 75 µm, Laufenberg) is applied as the peel-off film. The finished systems are punched out in sizes in the wanted form of from

5 to 30cm² using a punching tool. The complete systems are sealed individually in sachets of aluminised paper.

Example 3: Vial containing 1.0 mg dry, lyophilized 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid (mixed sodium salts thereof). After dilution with 1 ml of water, a solution (concentration 1 mg/ml) for i.v. infusion is obtained.

Composition:

active ingredient (free diphosphonic acid)		1.0 mg
mannitol		46.0 mg
Trisodium citrate x 2 H ₂ O	ca.	3.0 mg
water		1 ml
water for injection		1 ml

In 1 ml of water, the active ingredient is titrated with trisodium citrate x 2 H₂O to pH 6.0. Then, the mannitol is added and the solution is lyophilized and the lyophilisate filled into a vial.

Example 4: Ampoule containing active ingredient, for instance disodium pamidronate pentahydrate dissolved in water. The solution (concentration 3 mg/ml) is for i.v. infusion after dilution.

Composition:

active ingredient	19.73 mg
(Δ 5.0 mg of anhydrous active ingredient)	
mannitol	250 mg
water for injection	5 ml

Example 5 Treatment of Patients

“A Multinational, Randomized, Double-Blind, Placebo-Controlled, Parallel-Group, Dose-Ranging, Safety and Efficacy Trial With Intravenous Bolus Injections of Zoledronate In the Treatment of Postmenopausal Osteoporosis”

This was a dose and dose regimen-finding 12 month trial of iv zoledronic acid in patients with postmenopausal osteoporosis. Three hundred and fifty one patients were randomized to six study arms. Patients who had recent exposure to bone active drugs, e.g. bisphosphonates, estrogen, calcitonin, raloxifene, or a history of metabolic bone diseases were excluded. All patients were evaluated at baseline and in 3-monthly visits. Zoledronic acid or placebo was administered as a bolus iv injection into a peripheral vein over 5 minutes at every visit.

Efficacy was ascertained by measurement of percent change from baseline in bone mineral density (BMD) measured by dual energy X-ray absorptiometry (DEXA) as compared to placebo, at 6, 9, and 12 months.

As a special safety measure trans-iliac bone biopsies were obtained in a subset of patients from all study arms at 12 months, and X-rays of the thoracic and lumbar spine from all study participants were evaluated at baseline and at 12 months for the occurrence of incident vertebral fractures.

Additionally, the degree and duration of suppression of biochemical markers of bone turnover - parathyroid hormone (PTH), bone specific alkaline phosphatase (BSAP), serum C-telopeptide (CTX), serum osteocalcin, urine N-telopeptide (NTX)/creatinine ratio, urine deoxypyridinoline (d-pyd)/creatinine ratio, urine pyridinoline (pyd)/creatinine ratio - was obtained every 3 months and measured in a central laboratory.

Study Arms

- Placebo
- 0.25 mg zoledronic acid every 3 months
- 0.5 mg zoledronic acid every 3 months
- 1.0 mg zoledronic acid every 3 months

- 2.0 mg zoledronic acid every 6 months
- 4.0 mg zoledronic acid every 12 months

The 12 month results showed that all treatment arms demonstrated a percent change from baseline in BMD significantly ($p < 0.001$) greater than placebo and not dissimilar one from another.

Summary of stepwise multiple comparisons of the active doses of zoledronate versus placebo for percent change from baseline in bone mineral density of the lumbar spine; postero anterior (L1-L4) at 12 months Confirmatory analysis ITT population					
Step Number	Most significant contrast	Difference	Standard error of difference	Lower 97.5% confidence limit	p-value
1	zoledronate 4 x 0.25 mg - placebo	5.1	0.55	3.7	<0.001
2	zoledronate 4 x 0.5 mg - placebo	4.9	0.56	3.5	<0.001
3	zoledronate 1 x 4.0 mg - placebo	4.6	0.53	3.3	<0.001
4	zoledronate 4 x 1.0 mg - placebo	4.5	0.55	3.2	<0.001
5	zoledronate 2 x 2.0 mg - placebo	4.2	0.57	3.1	<0.001

Note: Stepwise multiple comparison of the active doses of zoledronate versus placebo at a one-sided multiple alpha level of 2.5% adjusting for multiple comparisons according to Marcus, Peritz and Gabriel (1976)

Bone mineral density increased compared to placebo at the spine, hip, distal radius, and "total body". Suppression of biochemical markers of bone formation and bone resorption confirmed and supported the BMD results, demonstrating suppression of bone turnover to the pre-menopausal level throughout the 6 and 12 month dosing intervals.

The BMD data indicate that zoledronic acid dose administration as infrequent as every 6 or 12 months can safely result in a statistically significant and medically relevant bone mass increase. It is believed that these data further indicate that a continued preservation of new bone beyond one year, without additional dose administration, is likely or that further bone mass increase is possible. It is also believed that re-treatment in additional cycles of every 6 month, 12 month, or less frequent dose administration will lead to further BMD increase. A reduction of risk of osteoporotic fracture is expected to accompany the bone mass increases.

CLAIMS

1. A method for the treatment of conditions of abnormally increased bone turnover in a patient in need of such treatment which comprises intermittently administering an effective amount of a bisphosphonate to the patient, wherein the period between administrations of bisphosphonate is at least about 6 months.
2. Use of a bisphosphonate in the preparation of a medicament for the treatment of conditions of abnormally increased bone turnover in which the bisphosphonate is administered intermittently and in which the period between administrations of bisphosphonate is at least about 6 months.
3. A method according to claim 1 or a use according to claim 2, wherein the period between administrations of bisphosphonate is at least about once a year.
4. A method according to claim 1 or use according to claim 2, in which the bisphosphonate is an N-bisphosphonate or a pharmaceutically acceptable salt thereof, or any hydrate thereof.
5. A method according to claim 1 or use according to claim 2, for the prophylactic treatment of osteoporosis wherein the period between administrations of bisphosphonate is about once per year or less frequent.
6. A method according to claim 1 or use according to claim 2, in which the bisphosphonate is 1-hydroxy-2- (imidazol-1-yl)ethane-1,1-diphosphonic acid, or a pharmaceutically acceptable salt thereof, or any hydrate thereof.
7. A method according to claim 1 or use according to claim 2, for the prophylactic treatment of osteoporosis wherein the period between administrations of bisphosphonate is about once every 18 months, about once every two years or less frequently.

8. A method according to claim 1 or a use according to claim 2 wherein the bisphosphonates is administered parenterally.
9. A method according to claim 1 or use according to claim 3 wherein the bisphosphonates is administered intravenously, subcutaneously, intramuscularly, or transdermally.
10. A method according to claim 1 or a use according to claim 2 in which the bisphosphonate is selected from: 3-amino-1-hydroxypropane-1,1-diphosphonic acid (pamidronic acid), e.g. pamidronate (APD); 3-(N,N-dimethylamino)-1-hydroxypropane-1,1-diphosphonic acid, e.g. dimethyl-APD; 4-amino-1-hydroxybutane-1,1-diphosphonic acid (alendronic acid), e.g. alendronate; 1-hydroxy-ethidene-bisphosphonic acid, e.g. etidronate; 1-hydroxy-3-(methylpentylamino)-propylidene-bisphosphonic acid, (ibandronic acid), e.g. ibandronate; 6-amino-1-hydroxyhexane-1,1-diphosphonic acid, e.g. amino-hexyl-BP; 3-(N-methyl-N-pentylamino)-1-hydroxypropane-1,1-diphosphonic acid, e.g. methyl-pentyl-APD (= BM 21.0955); 1-hydroxy-2-(imidazol-1-yl)ethane-1,1-diphosphonic acid, e.g. zoledronic acid; 1-hydroxy-2-(3-pyridyl)ethane-1,1-diphosphonic acid (risedronic acid), e.g. risedronate, including N-methyl pyridinium salts thereof, for example N-methyl pyridinium iodides such as NE-10244 or NE-10446; 1-(4-chlorophenylthio)methane-1,1-diphosphonic acid (tiludronic acid), e.g. tiludronate; 3-[N-(2-phenylthioethyl)-N-methylamino]-1-hydroxypropane-1,1-diphosphonic acid; 1-hydroxy-3-(pyrrolidin-1-yl)propane-1,1-diphosphonic acid, e.g. EB 1053 (Leo); 1-(N-phenylaminothiocarbonyl)methane-1,1-diphosphonic acid, e.g. FR 78844 (Fujisawa); 5-benzoyl-3,4-dihydro-2H-pyrazole-3,3-diphosphonic acid tetraethyl ester, e.g. U-81581 (Upjohn); 1-hydroxy-2-(imidazo[1,2-a]pyridin-3-yl)ethane-1,1-diphosphonic acid, e.g. YM 529; and 1,1-dichloromethane-1,1-diphosphonic acid (clodronic acid), e.g. clodronate; YM175, or a pharmaceutically acceptable salt thereof, or any hydrate thereof.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 December 2001 (27.12.2001)

PCT

(10) International Publication Number
WO 01/97788 A3

- (51) International Patent Classification: **A61K 31/663**, A61P 19/08, 19/10
- (21) International Application Number: **PCT/EP01/06850**
- (22) International Filing Date: **18 June 2001 (18.06.2001)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
09/597,135 20 June 2000 (20.06.2000) US
60/267,689 9 February 2001 (09.02.2001) US
- (71) Applicant (for all designated States except AT, US): **NOVARTIS AG [CH/CH]**; Lichtstrasse 35, CH-4056 Basel (CH).
- (71) Applicant (for AT only): **NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT M.B.H. [AT/AT]**; Brunner Strasse 59, A-1230 Vienna (AT).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **HOROWITZ, Zebulun, D.** [US/US]; 83 Blackburn Road, Basking Ridge, NJ 07920 (US). **RICHARDSON, Peter, C.** [GB/US]; 4 Spring Lake Drive, Far Hills, NJ 07931 (US). **TRECHSEL, Ulrich** [CH/CH]; Grosshaus, CH-7242 Luzein (CH).
- (54) Agent: **BECKER, Konrad**; Novartis AG, Corporate Intellectual Property, Patent & Trademark Department, CH-4002 Basel (CH).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published: **with international search report**
- (88) Date of publication of the international search report: **27 June 2002**
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/97788 A3

(54) Title: **METHOD OF ADMINISTERING BISPHOSPHONATES**

(57) Abstract: Bisphosphonates, in particular more potent N-bisphosphonates such as zoledronic acid and derivatives, can be used with satisfactory results for prolonged inhibition of bone resorption in conditions of abnormally increased bone turnover, e.g. osteoporosis, by intermittent administration, wherein the periods between bisphosphonate administrations are longer than was previously considered appropriate, e.g. a dosing interval of at least about 6 months or less frequently.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/06850

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/663 A61P19/08 A61P19/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 6 015 801 A (SANTORA II ARTHUR C ET AL) 18 January 2000 (2000-01-18) column 6, line 63 -column 14, line 45; claims 1-44,46-58; examples 2-7	2-10
Y	WO 99 18972 A (ASTRA AB ;UNGELL ANNA LENA (SE)) 22 April 1999 (1999-04-22) claims 1-20; examples 2-4	2-10
P,Y	WO 00 71104 A (NOVARTIS ERFIND VERWALT GMBH ;NOVARTIS AG (CH); GREEN JONATHAN (CH) 30 November 2000 (2000-11-30) page 1, line 1 -page 11, line 31; claims 1-10; examples 1-7	2-10
Y	DE 198 53 483 A (PRODOTTI ANTIBIOTICI SPA) 27 May 1999 (1999-05-27) claims 1-10; examples 1-8	2-10
	-/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

26 March 2002

Date of mailing of the international search report

09/04/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kling, I

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/06850

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
------------	--	-----------------------

Y	<p>WO 99 15155 A (RANTALA PERTTI TAPANI ;LEIRAS OY (FI); LEHTOLA VELI MATTI (FI); RA) 1 April 1999 (1999-04-01) page 1, line 4 -page 3, line 28; claims 1-6</p> <p>-----</p>	2-10
---	--	------

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/06850

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 6015801	A	18-01-2000	GB 2336311 A	20-10-1999
			AT 209037 T	15-12-2001
			AU 8493698 A	16-02-1999
			BG 104093 A	31-10-2000
			BR 9810779 A	25-07-2000
			CN 1265035 T	30-08-2000
			DE 69803291 D1	21-02-2002
			EE 200000040 A	16-10-2000
			EP 1132088 A2	12-09-2001
			EP 1151752 A2	07-11-2001
			EP 1175903 A2	30-01-2002
			EP 1175904 A2	30-01-2002
			EP 0998292 A2	10-05-2000
			ES 2164447 T3	16-02-2002
			HR 20000035 A1	28-02-2001
			HU 0004653 A2	28-04-2001
			JP 2002502433 T	22-01-2002
			NO 20000323 A	20-03-2000
			PL 338635 A1	06-11-2000
			SK 932000 A3	12-09-2000
			TR 200000145 T2	22-05-2000
			WO 9904773 A2	04-02-1999
			US 5994329 A	30-11-1999
			US 6333316 B1	25-12-2001
			US 2002010161 A1	24-01-2002
			US 6225294 B1	01-05-2001
			US 2001001306 A1	17-05-2001
			US 2001002395 A1	31-05-2001
			US 2001018431 A1	30-08-2001
			ZA 9806479 A	22-01-1999
WO 9918972	A	22-04-1999	AU 9469198 A	03-05-1999
			EP 1021194 A1	26-07-2000
			JP 2001519398 T	23-10-2001
			WO 9918972 A1	22-04-1999
			ZA 9808886 A	12-04-1999
WO 0071104	A	30-11-2000	AU 5214100 A	12-12-2000
			WO 0071104 A2	30-11-2000
			EP 1178810 A2	13-02-2002
			NO 20015638 A	15-01-2002
DE 19853483	A	27-05-1999	IT 1296495 B1	25-06-1999
			BE 1012137 A3	02-05-2000
			DE 19853483 A1	27-05-1999
			ES 2152865 A1	01-02-2001
			FR 2771638 A1	04-06-1999
			GB 2331459 A	26-05-1999
WO 9915155	A	01-04-1999	FI 973733 A	20-03-1999
			AU 737738 B2	30-08-2001
			AU 9268598 A	12-04-1999
			BG 104265 A	30-11-2000
			BR 9812480 A	26-09-2000
			CA 2301185 A1	01-04-1999
			CN 1270516 T	18-10-2000
			EE 200000096 A	15-12-2000
			EP 1027037 A1	16-08-2000

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/06850

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9915155	A	WO 9915155 A1	01-04-1999
		HR 20000142 A1	30-06-2000
		HU 0004383 A2	28-08-2001
		JP 2001517616 T	09-10-2001
		NO 20001417 A	19-05-2000
		PL 339290 A1	04-12-2000
		SK 3932000 A3	07-11-2000
		TR 200000722 T2	21-11-2000